

# Uniqueness of the Fisher–Rao metric on the space of smooth densities

Peter W. Michor

University of Vienna, Austria  
[www.mat.univie.ac.at/~michor](http://www.mat.univie.ac.at/~michor)

IGAIA IV

Information Geometry and its Applications IV

June 12-17, 2016, Liblice, Czech Republic

In honor of Shun-ichi Amari

## Based on:

[M.Bauer, M.Bruveris, P.Michor: Uniqueness of the Fisher–Rao metric on the space of smooth densities, Bull. London Math. Soc. doi:10.1112/blms/bdw020]

[M.Bruveris, P.Michor: Geometry of the Fisher-Rao metric on the space of smooth densities]

[M.Bruveris, P. Michor, A.Parusinski, A. Rainer: Moser's Theorem for manifolds with corners, arxiv:1604.07787]

[M.Bruveris,P.Michor, A.Rainer: Determination of all diffeomorphism invariant tensor fields on the space of smooth positive densities on a compact manifold with corners]

The infinite dimensional geometry used here is based on:

[Andreas Kriegl, Peter W. Michor: The Convenient Setting of Global Analysis. Mathematical Surveys and Monographs, Volume: 53, Amer. Math. Soc., 1997]

Wikipedia [[https://en.wikipedia.org/wiki/Convenient\\_vector\\_space](https://en.wikipedia.org/wiki/Convenient_vector_space)]

# Abstract

For a smooth compact manifold  $M$ , any weak Riemannian metric on the space of smooth positive densities which is invariant under the right action of the diffeomorphism group  $Diff(M)$  is of the form

$$G_\mu(\alpha, \beta) = C_1(\mu(M)) \int_M \frac{\alpha \beta}{\mu} \mu + C_2(\mu(M)) \int_M \alpha \cdot \int_M \beta$$

for smooth functions  $C_1, C_2$  of the total volume  $\mu(M) = \int_M \mu$ .

In this talk the result is extended to:

- (0) Geometry of the Fisher-Rao metric: geodesics and curvature.
- (1) manifolds with boundary, for manifolds with corner.
- (2) to tensor fields of the form  $G_\mu(\alpha_1, \alpha_2, \dots, \alpha_k)$  for any  $k$  which are invariant under  $Diff(M)$ .

The Fisher–Rao metric on the space  $\text{Prob}(M)$  of probability densities is of importance in the field of information geometry. Restricted to finite-dimensional submanifolds of  $\text{Prob}(M)$ , so-called statistical manifolds, it is called Fisher’s information metric [Amari: Differential-geometrical methods in statistics, 1985]. The Fisher–Rao metric is invariant under the action of the diffeomorphism group. A uniqueness result was established [Čencov: Statistical decision rules and optimal inference, 1982, p. 156] for Fisher’s information metric on finite sample spaces and [Ay, Jost, Le, Schwachhöfer, 2014] extended it to infinite sample spaces.

The Fisher–Rao metric on the infinite-dimensional manifold of all positive probability densities was studied in [Friedrich: Die Fisher-Information und symplektische Strukturen, 1991], including the computation of its curvature.

# The space of densities

Let  $M^m$  be a smooth manifold. Let  $(U_\alpha, u_\alpha)$  be a smooth atlas for it. The *volume bundle*  $(\text{Vol}(M), \pi_M, M)$  of  $M$  is the 1-dimensional vector bundle (line bundle) which is given by the following cocycle of transition functions:

$$\psi_{\alpha\beta} : U_{\alpha\beta} = U_\alpha \cap U_\beta \rightarrow \mathbb{R} \setminus \{0\} = GL(1, \mathbb{R}),$$

$$\psi_{\alpha\beta}(x) = |\det d(u_\beta \circ u_\alpha^{-1})(u_\alpha(x))| = \frac{1}{|\det d(u_\alpha \circ u_\beta^{-1})(u_\beta(x))|}.$$

$\text{Vol}(M)$  is a trivial line bundle over  $M$ . But there is no natural trivialization. There is a natural order on each fiber. Since  $\text{Vol}(M)$  is a natural bundle of order 1 on  $M$ , there is a natural action of the group  $\text{Diff}(M)$  on  $\text{Vol}(M)$ , given by

$$\begin{array}{ccc} \text{Vol}(M) & \xrightarrow{|\det(T\varphi^{-1})| \circ \varphi} & \text{Vol}(M) \\ \downarrow & & \downarrow \\ M & \xrightarrow{\varphi} & M \end{array}$$

If  $M$  is orientable, then  $\text{Vol}(M) = \Lambda^m T^*M$ . If  $M$  is not orientable, let  $\tilde{M}$  be the orientable double cover of  $M$  with its deck-transformation  $\tau : \tilde{M} \rightarrow \tilde{M}$ . Then  $\Gamma(\text{Vol}(M))$  is isomorphic to the space  $\{\omega \in \Omega^m(\tilde{M}) : \tau^*\omega = -\omega\}$ . These are the ‘formes impaires’ of de Rham. See [M 2008, 13.1] for this.

Sections of the line bundle  $\text{Vol}(M)$  are called densities. The space  $\Gamma(\text{Vol}(M))$  of all smooth sections is a Fréchet space in its natural topology; see [Kriegel-M, 1997]. For each section  $\alpha$  of  $\text{Vol}(M)$  of compact support the integral  $\int_M \alpha$  is invariantly defined as follows: Let  $(U_\alpha, u_\alpha)$  be an atlas on  $M$  with associated trivialization  $\psi_\alpha : \text{Vol}(M)|_{U_\alpha} \rightarrow \mathbb{R}$ , and let  $f_\alpha$  be a partition of unity with  $\text{supp}(f_\alpha) \subset U_\alpha$ . Then we put

$$\int_M \mu = \sum_\alpha \int_{U_\alpha} f_\alpha \mu := \sum_\alpha \int_{u_\alpha(U_\alpha)} f_\alpha(u_\alpha^{-1}(y)) \cdot \psi_\alpha(\mu(u_\alpha^{-1}(y))) dy.$$

The integral is independent of the choice of the atlas and the partition of unity.

# The Fisher–Rao metric

Let  $M^m$  be a smooth compact manifold without boundary. Let  $\text{Dens}_+(M)$  be the space of smooth positive densities on  $M$ , i.e.,  $\text{Dens}_+(M) = \{\mu \in \Gamma(\text{Vol}(M)) : \mu(x) > 0 \forall x \in M\}$ .

Let  $\text{Prob}(M)$  be the subspace of positive densities with integral 1.

For  $\mu \in \text{Dens}_+(M)$  we have  $T_\mu \text{Dens}_+(M) = \Gamma(\text{Vol}(M))$  and for  $\mu \in \text{Prob}(M)$  we have

$T_\mu \text{Prob}(M) = \{\alpha \in \Gamma(\text{Vol}(M)) : \int_M \alpha = 0\}$ .

The Fisher–Rao metric on  $\text{Prob}(M)$  is defined as:

$$G_\mu^{\text{FR}}(\alpha, \beta) = \int_M \frac{\alpha}{\mu} \frac{\beta}{\mu} \mu.$$

It is invariant for the action of  $\text{Diff}(M)$  on  $\text{Prob}(M)$ :

$$\begin{aligned} \left( (\varphi^*)^* G_\mu^{\text{FR}} \right)_\mu (\alpha, \beta) &= G_{\varphi^* \mu}^{\text{FR}}(\varphi^* \alpha, \varphi^* \beta) = \\ &= \int_M \left( \frac{\alpha}{\mu} \circ \varphi \right) \left( \frac{\beta}{\mu} \circ \varphi \right) \varphi^* \mu = \int_M \frac{\alpha}{\mu} \frac{\beta}{\mu} \mu. \end{aligned}$$

## Theorem [BBM, 2016]

Let  $M$  be a compact manifold without boundary of dimension  $\geq 2$ . Let  $G$  be a smooth (equivalently, bounded) bilinear form on  $\text{Dens}_+(M)$  which is invariant under the action of  $\text{Diff}(M)$ . Then

$$G_\mu(\alpha, \beta) = C_1(\mu(M)) \int_M \frac{\alpha \beta}{\mu \mu} \mu + C_2(\mu(M)) \int_M \alpha \cdot \int_M \beta$$

for smooth functions  $C_1, C_2$  of the total volume  $\mu(M)$ .

To see that this theorem implies the uniqueness of the Fisher–Rao metric, note that if  $G$  is a  $\text{Diff}(M)$ -invariant Riemannian metric on  $\text{Prob}(M)$ , then we can equivariantly extend it to  $\text{Dens}_+(M)$  via

$$G_\mu(\alpha, \beta) = G_{\frac{\mu}{\mu(M)}} \left( \alpha - \left( \int_M \alpha \right) \frac{\mu}{\mu(M)}, \beta - \left( \int_M \beta \right) \frac{\mu}{\mu(M)} \right).$$



# Relations to right-invariant metrics on diffeom. groups

Let  $\mu_0 \in \text{Prob}(M)$  be a fixed smooth probability density. In [Khesin, Lenells, Misiolek, Preston, 2013] it has been shown, that the degenerate,  $\dot{H}^1$ -metric  $\frac{1}{2} \int_M \text{div}^{\mu_0}(X) \cdot \text{div}^{\mu_0}(X) \cdot \mu_0$  on  $\mathfrak{X}(M)$  is invariant under the adjoint action of  $\text{Diff}(M, \mu_0)$ . Thus the induced degenerate right invariant metric on  $\text{Diff}(M)$  descends to a metric on  $\text{Prob}(M) \cong \text{Diff}(M, \mu_0) \backslash \text{Diff}(M)$  via

$$\text{Diff}(M) \ni \varphi \mapsto \varphi^* \mu_0 \in \text{Prob}(M)$$

which is invariant under the right action of  $\text{Diff}(M)$ . This is the Fisher–Rao metric on  $\text{Prob}(M)$ . In [Modin, 2014], the  $\dot{H}^1$ -metric was extended to a non-degenerate metric on  $\text{Diff}(M)$ , also descending to the Fisher–Rao metric.

**Corollary.** *Let  $\dim(M) \geq 2$ . If a weak right-invariant (possibly degenerate) Riemannian metric  $\tilde{G}$  on  $\text{Diff}(M)$  descends to a metric  $G$  on  $\text{Prob}(M)$  via the right action, i.e., the mapping  $\varphi \mapsto \varphi^* \mu_0$  from  $(\text{Diff}(M), \tilde{G})$  to  $(\text{Prob}(M), G)$  is a Riemannian submersion, then  $G$  has to be a multiple of the Fisher–Rao metric.*

Note that any right invariant metric  $\tilde{G}$  on  $\text{Diff}(M)$  descends to a metric on  $\text{Prob}(M)$  via  $\varphi \mapsto \varphi_* \mu_0$ ; but this is not  $\text{Diff}(M)$ -invariant in general.

## Invariant metrics on $\text{Dens}_+(S^1)$ .

$\text{Dens}_+(S^1) = \Omega_+^1(S^1)$ , and  $\text{Dens}_+(S^1)$  is  $\text{Diff}(S^1)$ -equivariantly isomorphic to the space of all Riemannian metrics on  $S^1$  via  $\Phi = (\ )^2 : \text{Dens}_+(S^1) \rightarrow \text{Met}(S^1)$ ,  $\Phi(fd\theta) = f^2d\theta^2$ .

On  $\text{Met}(S^1)$  there are many  $\text{Diff}(S^1)$ -invariant metrics; see [Bauer, Harms, M, 2013]. For example Sobolev-type metrics. Write  $g \in \text{Met}(S^1)$  in the form  $g = \tilde{g}d\theta^2$  and  $h = \tilde{h}d\theta^2$ ,  $k = \tilde{k}d\theta^2$  with  $\tilde{g}, \tilde{h}, \tilde{k} \in C^\infty(S^1)$ . The following metrics are  $\text{Diff}(S^1)$ -invariant:

$$G_g^l(h, k) = \int_{S^1} \frac{\tilde{h}}{\tilde{g}} \cdot (1 + \Delta^g)^n \left( \frac{\tilde{k}}{\tilde{g}} \right) \sqrt{\tilde{g}} d\theta;$$

here  $\Delta^g$  is the Laplacian on  $S^1$  with respect to the metric  $g$ . The pullback by  $\Phi$  yields a  $\text{Diff}(S^1)$ -invariant metric on  $\text{Dens}_+(M)$ :

$$G_\mu(\alpha, \beta) = 4 \int_{S^1} \frac{\alpha}{\mu} \cdot \left( 1 + \Delta^{\Phi(\mu)} \right)^n \left( \frac{\beta}{\mu} \right) \mu.$$

For  $n = 0$  this is 4 times the Fisher–Rao metric. For  $n \geq 1$  we get different  $\text{Diff}(S^1)$ -invariant metrics on  $\text{Dens}_+(M)$  and on  $\text{Prob}(S^1)$ .

# Main Theorem

Let  $M$  be a compact manifold, possibly with corners, of dimension  $\geq 2$ . Let  $G$  be a smooth (equivalently, bounded)  $\binom{0}{n}$ -tensor field on  $\text{Dens}_+(M)$  which is invariant under the action of  $\text{Diff}(M)$ . If  $M$  is not orientable or if  $n \leq \dim(M) = m$ , then

$$\begin{aligned} G_\mu(\alpha_1, \dots, \alpha_n) &= C_0(\mu(M)) \int_M \frac{\alpha_1}{\mu} \dots \frac{\alpha_n}{\mu} \mu \\ &+ \sum_{i=1}^n C_i(\mu(M)) \int_M \alpha_i \cdot \int_M \frac{\alpha_1}{\mu} \dots \frac{\widehat{\alpha}_i}{\mu} \dots \frac{\alpha_n}{\mu} \mu \\ &+ \sum_{i < j}^n C_{ij}(\mu(M)) \int_M \frac{\alpha_i}{\mu} \frac{\alpha_j}{\mu} \mu \cdot \int_M \frac{\alpha_1}{\mu} \dots \frac{\widehat{\alpha}_i}{\mu} \dots \frac{\widehat{\alpha}_j}{\mu} \dots \frac{\alpha_n}{\mu} \mu \\ &+ \dots \\ &+ C_{12\dots n}(\mu(M)) \int_M \frac{\alpha_1}{\mu} \mu \cdot \int_M \frac{\alpha_2}{\mu} \mu \cdot \dots \int_M \frac{\alpha_n}{\mu} \mu. \end{aligned}$$

for some smooth functions  $C_0, \dots$  of the total volume  $\mu(M)$ .

## Main Theorem, continued

If  $M$  is orientable and  $n > \dim(M) = m$ , then each integral over more than  $m$  functions  $\alpha_i/\mu$  has to be replaced by the following expression which we write only for the first term:

$$C_0(\mu(M)) \int_M \frac{\alpha_1}{\mu} \cdots \frac{\alpha_n}{\mu} \mu + \\ + \sum C_0^K(\mu(M)) \int \frac{\alpha_{k_1}}{\mu} \cdots \frac{\alpha_{k_{n-m}}}{\mu} d\left(\frac{\alpha_{k_{n-m+1}}}{\mu}\right) \wedge \cdots \wedge d\left(\frac{\alpha_{k_n}}{\mu}\right)$$

where  $K = \{k_{n-m+1}, \dots, k_n\}$  runs through all subsets of  $\{1, \dots, n\}$  containing exactly  $m$  elements.

# Moser's theorem for manifolds with corners

[BMPR16]

*Let  $M$  be a compact smooth manifold with corners, possibly non-orientable. Let  $\mu_0$  and  $\mu_1$  be two smooth positive densities in  $\text{Dens}_+(M)$  with  $\int_M \mu_0 = \int_M \mu_1$ . Then there exists a diffeomorphism  $\varphi : M \rightarrow M$  such that  $\mu_1 = \varphi^* \mu_0$ . If and only if  $\mu_0(x) = \mu_1(x)$  for each corner  $x \in \partial^{\geq 2} M$  of codimension  $\geq 2$ , then  $\varphi$  can be chosen to be the identity on  $\partial M$ .*

This result is highly desirable even for  $M$  a simplex. The proof is essentially contained in [Banyaga1974], who proved it for manifolds with boundary.

# Geometry of the Fisher-Rao metric

$$G_\mu(\alpha, \beta) = C_1(\mu(M)) \int_M \frac{\alpha \beta}{\mu} \mu + C_2(\mu(M)) \int_M \alpha \cdot \int_M \beta$$

This metric will be studied in different representations.

$$\text{Dens}_+(M) \xrightarrow{R} C^\infty(M, \mathbb{R}_{>0}) \xrightarrow{\Phi} \mathbb{R}_{>0} \times S \cap C^\infty_{>0} \xrightarrow{W \times \text{Id}} (W_-, W_+) \times S \cap C^\infty_{>0}.$$

We fix  $\mu_0 \in \text{Prob}(M)$  and consider the mapping

$$R : \text{Dens}_+(M) \rightarrow C^\infty(M, \mathbb{R}_{>0}), \quad R(\mu) = f = \sqrt{\frac{\mu}{\mu_0}}.$$

The map  $R$  is a diffeomorphism and we will denote the induced metric by  $\tilde{G} = (R^{-1})^* G$ ; it is given by the formula

$$\tilde{G}_f(h, k) = 4C_1(\|f\|^2) \langle h, k \rangle + 4C_2(\|f\|^2) \langle f, h \rangle \langle f, k \rangle,$$

and this formula makes sense for  $f \in C^\infty(M, \mathbb{R}) \setminus \{0\}$ .

The map  $R$  is inspired by [B. Khesin, J. Lenells, G. Misiolek, S. C.

Preston: Geometry of diffeomorphism groups, complete integrability and geometric statistics. *Geom. Funct. Anal.*, 23(1):334–366, 2013.]

## Remark on $R^{-1}$

$$R^{-1} : C^\infty(M, \mathbb{R}) \rightarrow \Gamma_{\geq 0}(\text{Vol}(M)), \quad f \mapsto f^2 \mu_0$$

makes sense on the whole space  $C^\infty(M, \mathbb{R})$  and its image is stratified (loosely speaking) according to the rank of  $TR^{-1}$ . The image looks somewhat like the orbit space of a discrete reflection group. Geodesics are mapped to curves which are geodesics in the interior  $\Gamma_{>0}(\text{Vol}(M))$ , and they are reflected following Snell's law at some hyperplanes in the boundary.



# Polar coordinates

on the pre-Hilbert space  $(C^\infty(M, \mathbb{R}), \langle \cdot, \cdot \rangle_{L^2(\mu_0)})$ . Let  $S = \{\varphi \in L^2(M, \mathbb{R}) : \int_M \varphi^2 \mu_0 = 1\}$  denote the  $L^2$ -sphere. Then

$$\Phi : C^\infty(M, \mathbb{R}) \setminus \{0\} \rightarrow \mathbb{R}_{>0} \times (S \cap C^\infty), \quad \Phi(f) = (r, \varphi) = \left( \|f\|, \frac{f}{\|f\|} \right)$$

is a diffeomorphism. We set  $\bar{G} = (\Phi^{-1})^* \tilde{G}$ ; the metric has the expression

$$\bar{G}_{r,\varphi} = g_1(r) \langle d\varphi, d\varphi \rangle + g_2(r) dr^2,$$

with  $g_1(r) = 4C_1(r^2)r^2$  and  $g_2(r) = 4(C_1(r^2) + C_2(r^2)r^2)$ . Finally we change the coordinate  $r$  diffeomorphically to

$$s = W(r) = 2 \int_1^r \sqrt{g_2(\rho)} d\rho.$$

Then, defining  $a(s) = 4C_1(r(s)^2)r(s)^2$ , we have

$$\bar{G}_{s,\varphi} = a(s) \langle d\varphi, d\varphi \rangle + ds^2.$$

Let  $W_- = \lim_{r \rightarrow 0^+} W(r)$  and  $W_+ = \lim_{r \rightarrow \infty} W(r)$ . Then  $W : \mathbb{R}_{>0} \rightarrow (W_-, W_+)$  is a diffeomorphism.

This completes the first row in Fig. 1.

$$\begin{array}{ccccccc}
 \text{Dens}_+(M) & \xrightarrow{R} & C^\infty(M, \mathbb{R}_{>0}) & \xrightarrow{\phi} & \mathbb{R}_{>0} \times S \cap C^\infty_{>0} & \xrightarrow{W \times \text{Id}} & (W_-, W_+) \times S \cap C^\infty_{>0} \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \text{Dens}(M) \setminus \{0\} & \xrightarrow{R} & C^0(M, \mathbb{R}) \setminus \{0\} & \xrightarrow{\phi} & \mathbb{R}_{>0} \times S \cap C^0 & \xrightarrow{W \times \text{Id}} & \mathbb{R} \times S \cap C^0 \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \Gamma_{L^1}(\text{Vol}(M)) \setminus \{0\} & \xrightarrow{R} & L^2(M, \mathbb{R}) \setminus \{0\} & \xrightarrow{\phi} & \mathbb{R}_{>0} \times S & \xrightarrow{W \times \text{Id}} & \mathbb{R} \times S
 \end{array}$$

**Figure:** Representations of  $\text{Dens}_+(M)$  and its completions. In the second and third rows we assume that  $(W_-, W_+) = (-\infty, +\infty)$  and we note that  $R$  is a diffeomorphism only in the first row.

Geodesic equation:

$$\begin{aligned}
 \nabla_{\partial_t}^S \varphi_t &= \partial_t (\log g_1(r)) \varphi_t \\
 r_{tt} &= \frac{C_0^2}{2} \frac{g_1'(r)}{g_1(r)^2 g_2(r)} - \frac{1}{2} \partial_t (\log g_2(r)) r_t
 \end{aligned}$$

Since  $\bar{G}$  induces the canonical metric on  $(W_-, W_+)$ , a necessary condition for  $\bar{G}$  to be complete is  $(W_-, W_+) = (-\infty, +\infty)$ .

Rewritten in terms of the functions  $C_1, C_2$  this becomes

$$W_+ = \infty \Leftrightarrow \left( \int_1^\infty r^{-1/2} \sqrt{C_1(r)} dr = \infty \text{ or } \int_1^\infty \sqrt{C_2(r)} dr = \infty \right),$$

and similarly for  $W_- = -\infty$ , with the limits of the integration being 0 and 1.

# Relation to hypersurfaces of revolution in the (pre-) Hilbert space

We consider the metric on  $(W_-, W_+) \times S \cap C^\infty$  in the form  $\tilde{G}_{r,\varphi} = a(s)\langle d\varphi, d\varphi \rangle + ds^2$  where  $a(s) = 4C_1(r(s)^2)r(s)^2$ . Then we consider the isometric embedding (remember  $\langle \varphi, d\varphi \rangle = 0$  on  $S \cap C^\infty$ )

$\Psi : ((W_-, W_+) \times S \cap C^\infty, \tilde{G}) \rightarrow (\mathbb{R} \times C^\infty(M, \mathbb{R}), du^2 + \langle df, df \rangle),$

$$\Psi(s, \varphi) = \left( \int_0^s \sqrt{1 - \frac{a'(\sigma)^2}{4a(\sigma)}} d\sigma, \sqrt{a(s)}\varphi \right),$$

which defined and smooth only on the open subset

$$R := \{(s, \varphi) \in (W_-, W_+) \times S \cap C^\infty : a'(s)^2 < 4a(s)\}.$$

Fix some  $\varphi_0 \in S \cap C^\infty$  and consider the generating curve

$$s \mapsto \left( \int_0^s \sqrt{1 - \frac{a'(\sigma)^2}{4a(\sigma)}} d\sigma, \sqrt{a(s)} \right) \in \mathbb{R}^2.$$

Then  $s$  is an arc-length parameterization of this curve!

Given any arc-length parameterized curve  $I \ni s \mapsto (c_1(s), c_2(s))$  in  $\mathbb{R}^2$  and its generated hypersurface of rotation

$$\{(c_1(s), c_2(s)\varphi) : s \in I, \varphi \in S \cap C^\infty\} \subset \mathbb{R} \times C^\infty(M, \mathbb{R}),$$

the induced metric in the  $(s, \varphi)$ -parameterization is  $ds^2 + c_2(s)^2 \langle d\varphi, d\varphi \rangle$ .

This suggests that the moduli space of hypersurfaces of revolution is naturally embedded in the moduli space of all metrics of the form (b).

# Theorem

*If  $(W_-, W_+) = (-\infty, +\infty)$ , then any two points  $(s_0, \varphi_0)$  and  $(s_1, \varphi_1)$  in  $\mathbb{R} \times S$  can be joined by a minimal geodesic. If  $\varphi_0$  and  $\varphi_1$  lie in  $S \cap C^\infty$ , then the minimal geodesic lies in  $\mathbb{R} \times S \cap C^\infty$ .*

**Proof.** If  $\varphi_0$  and  $\varphi_1$  are linearly independent, we consider the 2-space  $V = V(\varphi_0, \varphi_1)$  spanned by  $\varphi_0$  and  $\varphi_1$  in  $L^2$ . Then  $\mathbb{R} \times V \cap S$  is totally geodesic since it is the fixed point set of the isometry  $(s, \varphi) \mapsto (s, \mathfrak{s}_V(\varphi))$  where  $\mathfrak{s}_V$  is the orthogonal reflection at  $V$ . Thus there exists a minimizing geodesic between  $(s_0, \varphi_0)$  and  $(s_1, \varphi_1)$  in the complete 3-dimensional Riemannian submanifold  $\mathbb{R} \times V \cap S$ . This geodesic is also length-minimizing in the strong Hilbert manifold  $\mathbb{R} \times S$  by the following arguments:

Given any smooth curve  $c = (s, \varphi) : [0, 1] \rightarrow \mathbb{R} \times S$  between these two points, there is a subdivision  $0 = t_0 < t_1 < \dots < t_N = 1$  such that the piecewise geodesic  $c_1$  which first runs along a geodesic from  $c(t_0)$  to  $c(t_1)$ , then to  $c(t_2)$ ,  $\dots$ , and finally to  $c(t_N)$ , has length  $\text{Len}(c_1) \leq \text{Len}(c)$ . This piecewise geodesic now lies in the totally geodesic  $(N + 2)$ -dimensional submanifold  $\mathbb{R} \times V(\varphi(t_0), \dots, \varphi(t_N)) \cap S$ . Thus there exists a geodesic  $c_2$  between the two points  $(s_0, \varphi_0)$  and  $(s_1, \varphi_1)$  which is length minimizing in this  $(N + 2)$ -dimensional submanifold. Therefore  $\text{Len}(c_2) \leq \text{Len}(c_1) \leq \text{Len}(c)$ . Moreover,  $c_2 = (s \circ c_2, \varphi \circ c_2)$  lies in  $\mathbb{R} \times V(\varphi_0, (\varphi \circ c_2)'(0)) \cap S$  which also contains  $\varphi_1$ , thus  $c_2$  lies in  $\mathbb{R} \times V(\varphi_0, \varphi_1) \cap S$ .

If  $\varphi_0 = \varphi_1$ , then  $\mathbb{R} \times \{\varphi_0\}$  is a minimal geodesic. If  $\varphi_0 = -\varphi_0$  we choose a great circle between them which lies in a 2-space  $V$  and proceed as above. □

# Covariant derivative

On  $\mathbb{R} \times S$  (we assume that  $(W_-, W_+) = \mathbb{R}$ ) with metric  $\bar{G} = ds^2 + a(s)\langle d\varphi, d\varphi \rangle$  we consider smooth vector fields  $f(s, \varphi)\partial_s + X(s, \varphi)$  where  $X(s, \varphi) \in \mathfrak{X}(S)$  is a smooth vector field on the Hilbert sphere  $S$ . We denote by  $\nabla^S$  the covariant derivative on  $S$  and get

$$\begin{aligned} \nabla_{f\partial_s + X}(g\partial_s + Y) &= (f \cdot g_s + dg(X) - \frac{a_s}{2}\langle X, Y \rangle)\partial_s \\ &\quad + \frac{a_s}{2a}(fY + gX) + fY_s + \nabla_X^S Y \end{aligned}$$

Curvature:

$$\begin{aligned} \mathcal{R}(f\partial_s + X, g\partial_s + Y)(h\partial_s + Z) &= \\ &= \left(\frac{a_{ss}}{2} - \frac{a_s^2}{4a}\right)\langle gX - fY, Z \rangle\partial_s + \mathcal{R}^S(X, Y)Z \\ &\quad - \left(\left(\frac{a_s}{2a}\right)_s + \frac{a_s^2}{4a^2}\right)h(gX - fY) + \frac{a_s}{2a}(\langle X, Z \rangle Y - \langle Y, Z \rangle X). \end{aligned}$$



# Sectional Curvature

Let us take  $X, Y \in T_\varphi S$  with  $\langle X, Y \rangle = 0$  and  $\langle X, X \rangle = \langle Y, Y \rangle = 1/a(s)$ , then

$$\begin{aligned}\text{Sec}_{(s,\varphi)}(\text{span}(X, Y)) &= \frac{1}{a} - \frac{a_s}{2a^2}, \\ \text{Sec}_{(s,\varphi)}(\text{span}(\partial_s, Y)) &= -\frac{a_{ss}}{2a} + \frac{a_s^2}{4a^2}\end{aligned}$$

are all the possible sectional curvatures.

## Back to the Main Theorem

Let  $M$  be a compact manifold, possibly with corners, of dimension  $\geq 2$ . Then the space of all  $\text{Diff}(M)$ -invariant purely covariant tensor fields on  $\text{Dens}_+(M)$  is generated as algebra with unit 1 over the ring of smooth functions  $f(\mu(M))$ ,  $f \in C^\infty(\mathbb{R}, \mathbb{R})$  by the following generators, allowing for permutations of the entries  $\alpha_i \in T_\mu \text{Dens}_+(M)$ :

$$\int_M \frac{\alpha_1}{\mu} \cdots \frac{\alpha_n}{\mu} \mu \quad \text{for all } n \in \mathbb{N}_{>0}, \text{ and by}$$
$$\int \frac{\alpha_1}{\mu} \cdots \frac{\alpha_{n-m}}{\mu} d\left(\frac{\alpha_{n-m+1}}{\mu}\right) \wedge \cdots \wedge d\left(\frac{\alpha_n}{\mu}\right)$$

for  $n > \dim(M)$  and orientable  $M$ .

# Manifolds with corners alias quadrantic (orthantic) manifolds

For more information we refer to [DouadyHerault73], [Michor80], [Melrose96], etc. Let  $Q = Q^m = \mathbb{R}_{\geq 0}^m$  be the positive orthant or quadrant. By Whitney's extension theorem or Seeley's theorem, restriction  $C^\infty(\mathbb{R}^m) \rightarrow C^\infty(Q)$  is a surjective continuous linear mapping which admits a continuous linear section (extension mapping); so  $C^\infty(Q)$  is a direct summand in  $C^\infty(\mathbb{R}^m)$ . A point  $x \in Q$  is called a *corner of codimension*  $q > 0$  if  $x$  lies in the intersection of  $q$  distinct coordinate hyperplanes. Let  $\partial^q Q$  denote the set of all corners of codimension  $q$ .

A manifold with corners (recently also called a quadrantic manifold)  $M$  is a smooth manifold modelled on open subsets of  $Q^m$ . We assume that it is connected and second countable; then it is paracompact and for each open cover it admits a subordinated smooth partition of unity. Any manifold with corners  $M$  is a submanifold with corners of an open manifold  $\tilde{M}$  of the same dim. Restriction  $C^\infty(\tilde{M}) \rightarrow C^\infty(M)$  is a surjective continuous linear map which admits a continuous linear section. Thus  $C^\infty(M)$  is a topological direct summand in  $C^\infty(\tilde{M})$  and the same holds for the dual spaces: The space of distributions  $\mathcal{D}'(M)$ , which we identify with  $C^\infty(M)'$ , is a direct summand in  $\mathcal{D}'(\tilde{M})$ . It consists of all distributions with support in  $M$ .

We do not assume that  $M$  is oriented, but eventually we will assume that  $M$  is compact. Diffeomorphisms of  $M$  map the boundary  $\partial M$  to itself and map the boundary  $\partial^q M$  of corners of codimension  $q$  to itself;  $\partial^q M$  is a submanifold of codimension  $q$  in  $M$ ; in general  $\partial^q M$  has finitely many connected components. We shall consider  $\partial M$  as stratified into the connected components of all  $\partial^q M$  for  $q > 0$ .

# Beginning of the proof of the Main Theorem

Fix a basic probability density  $\mu_0$ . By Moser's theorem for manifolds with corners, for each  $\mu \in \text{Dens}_+(M)$  there exists a diffeomorphism  $\varphi_\mu \in \text{Diff}(M)$  with  $\varphi_\mu^* \mu = \mu(M) \mu_0 =: c \cdot \mu_0$  where  $c = \mu(M) = \int_M \mu > 0$ . Then

$$\begin{aligned} ((\varphi_\mu^*)^* G)_\mu(\alpha_1, \dots, \alpha_n) &= G_{\varphi_\mu^* \mu}(\varphi_\mu^* \alpha_1, \dots, \varphi_\mu^* \alpha_n) = \\ &= G_{c \cdot \mu_0}(\varphi_\mu^* \alpha_1, \dots, \varphi_\mu^* \alpha_n). \end{aligned}$$

Thus it suffices to show that for any  $c > 0$  we have

$$G_{c\mu_0}(\alpha_1, \dots, \alpha_n) = C_0(c) \cdot \int_M \frac{\alpha_1}{\mu_0} \dots \frac{\alpha_n}{\mu_0} \mu_0 + \dots$$

for some functions  $C_0, \dots$  of the total volume  $c = \mu(M)$ . Since  $c \mapsto c \cdot \mu_0$  is a smooth curve in  $\text{Dens}_+(M)$ , the functions  $C_0, \dots$  are then smooth in  $c$ . All  $k$ -linear forms are still invariant under the action of the group

$$\text{Diff}(M, c\mu_0) = \text{Diff}(M, \mu_0) = \{\psi \in \text{Diff}(M) : \psi^* \mu_0 = \mu_0\}.$$

The  $k$ -linear form

$$(T_{\mu_0} \text{Dens}_+(M))^k \ni (\alpha_1, \dots, \alpha_n) \mapsto G_{c\mu_0} \left( \frac{\alpha_1}{\mu_0} \mu_0, \dots, \frac{\alpha_n}{\mu_0} \mu_0 \right)$$

can be viewed as a bounded  $k$ -linear form

$$C^\infty(M)^k \ni (f_1, \dots, f_n) \mapsto G_c(f_1, \dots, f_n).$$

Using the Schwartz kernel theorem,  $\check{G}_c$  has a kernel  $\hat{G}_c$ , which is a distribution (generalized function) in

$$\begin{aligned} \mathcal{D}'(M^n) &\cong \mathcal{D}'(M) \bar{\otimes} \dots \bar{\otimes} \mathcal{D}'(M) = (C^\infty(M) \bar{\otimes} \dots \bar{\otimes} C^\infty(M))' \\ &\cong L(C^\infty(M^k), \mathcal{D}'(M^{n-k})). \end{aligned}$$

Note the defining relations

$$G_c(f_1, \dots, f_n) = \langle \check{G}_c(f_1, \dots, f_k), f_{k+1} \otimes \dots \otimes f_n \rangle = \langle \hat{G}_c, f_1 \otimes \dots \otimes f_n \rangle.$$

$\hat{G}_c$  is invariant under the diagonal action of  $\text{Diff}(M, \mu_0)$  on  $M^n$ .

The infinitesimal version of this invariance is:

$$\begin{aligned} 0 &= \langle \mathcal{L}_{X^{\text{diag}}} \hat{G}_c, f_1 \otimes \cdots \otimes f_n \rangle = -\langle \hat{G}_c, \mathcal{L}_{X^{\text{diag}}}(f_1 \otimes \cdots \otimes f_n) \rangle \\ &= -\sum_{i=1}^n \langle \hat{G}_c, f_1 \otimes \cdots \otimes \mathcal{L}_X f_i \otimes \cdots \otimes f_n \rangle \end{aligned}$$

$$X^{\text{diag}} = X \times 0 \times \dots \times 0 + 0 \times X \times 0 \times \dots \times 0 + \dots$$

for all  $X \in \mathfrak{X}(M, \mu_0)$ .

We will consider various (permuted versions) of the associated bounded mappings

$$\check{G}_c : C^\infty(M)^k \rightarrow (C^\infty(M)^{n-k})' = \mathcal{D}'(M^{n-k}).$$

We shall use the fixed density  $\mu_0 \in \text{Dens}_+(M)$  for the rest of this section. So we identify distributions on  $M^k$  with the dual space  $C^\infty(M^k)' =: \mathcal{D}'(M^k)$

# The Lie algebra of $\text{Diff}(M, \mu_0)$

For a fixed positive density  $\mu_0$  on  $M$ , the Lie algebra of  $\text{Diff}(M, \mu_0)$  which we will denote by  $\mathfrak{X}(M, \partial M, \mu_0)$ , is the subalgebra of vector fields which are tangent to each boundary stratum and which are divergence free:  $0 = \text{div}^{\mu_0}(X) := \frac{\mathcal{L}_X \mu_0}{\mu_0}$ . These are exactly the fields  $X$  such that for each good subset  $U$  (where each density can be identified with an  $m$ -form) the form  $\hat{l}_{\mu_0}(X)$  is a closed form in  $\Omega^{m-1}(U, \partial U)$ , and  $0 = \text{div}^{\mu_0}(X) := \frac{\mathcal{L}_X \mu_0}{\mu_0}$ .

Denote by  $\mathfrak{X}_{\text{exact}}(M, \partial M, \mu_0)$  the set (not a vector space) of 'exact' divergence free vector fields  $X = \hat{l}_{\mu_0}^{-1}(d\omega)$ , where  $\omega \in \Omega_c^{m-2}(U, \partial U)$  for a good subset  $U \subset M$ . They are automatically tangent to each boundary stratum since  $d\omega \in \Omega_c^{m-1}(U, \partial U)$ .



**Lemma** *If for  $f \in C^\infty(M)$  and a good set  $U \subseteq M$  we have  $(\mathcal{L}_X f)|_U = 0$  for all  $X \in \mathfrak{X}_{\text{exact}}(M, \partial M, \mu_0)$ , then  $f|_U$  is constant.*

**Lemma** *If for a distribution  $A \in \mathcal{D}'(M) = C^\infty(M)'$  and a connected open set  $U \subseteq M$  we have  $\mathcal{L}_X A|_U = 0$  for all  $X \in \mathfrak{X}_{\text{exact}}(M, \partial M, \mu_0)$ , then  $A|_U = C\mu_0|_U$  for some constant  $C$ , meaning  $\langle A, f \rangle = C \int_M f \mu_0$  for all  $f \in C_c^\infty(U)$ .*

This lemma proves the theorem for the case  $n = 1$ .

**Lemma** *Each operator*

$$\check{G}_c : C^\infty(M) \rightarrow C^\infty(M^{n-1})'$$

$$f_i \mapsto ((f_1, \dots, \hat{f}_i, \dots, f_n) \mapsto G_c(f_1, \dots, f_n))$$

*has the following property: If for  $f \in C^\infty(M)$  and a connected open  $U \subseteq M$  the restriction  $f|_U$  is constant, then  $\mathcal{L}_{X^{\text{diag}}}(\check{G}_c(f))|_{U^{n-1}} = 0$  for each exact vector field  $X \in \mathfrak{X}_{\text{exact}}(M, \partial M, \mu_0)$ .*

**Lemma** Let  $\hat{G}$  be an invariant distribution in  $\mathcal{D}'(M^n)$ . Then for each  $1 \leq i \leq n$  there exists an invariant distribution  $\hat{G}_i \in \mathcal{D}'(M^{n-1})$  such that the distribution

$$(f_1, \dots, f_n) \mapsto \hat{G}(f_1, \dots, f_n) - \hat{G}_i(f_1, \dots, \hat{f}_i, \dots, f_n) \cdot \int_M f_i \mu_0$$

has support in the set

$$D_i(M) = \{(x_1, \dots, x_n) : x_i = x_j \text{ for some } j \neq i\}.$$

**Lemma** There exists a constant  $C = C(c)$  such that the distribution  $\hat{G}_c - C\mu_0^{\otimes n}$  is supported on the union of all partial diagonals

$$D := \{(x_1, \dots, x_n) \in M^n : \text{for at least one pair } i \neq j \\ \text{we have equality: } x_i = x_j\}.$$

**Lemma** Let  $\hat{G} \in \mathcal{D}'(M^n)$  be a  $\text{Diff}(M, \mu_0)$ -invariant distribution, supported on the full diagonal  $\Delta(M) = \{(x_1, \dots, x_n) \in M^n : x_1 = \dots = x_n\} \subset M^n$ . If  $n \leq \dim(M)$  or if  $M$  is not orientable, there exist some constant  $C$  such that  $G(f_1, \dots, f_n) = C \int_M f_1 \dots f_n \mu_0$ . If  $n > \dim(M)$  and if  $M$  is orientable, then there exist constants such that

$$C_0 \int_M \frac{\alpha_1}{\mu} \dots \frac{\alpha_n}{\mu} \mu + \sum C_0^K \int \frac{\alpha_{k_1}}{\mu} \dots \frac{\alpha_{k_{n-m}}}{\mu} d\left(\frac{\alpha_{k_{n-m+1}}}{\mu}\right) \wedge \dots \wedge d\left(\frac{\alpha_{k_n}}{\mu}\right)$$

where  $K = \{k_{n-m+1}, \dots, k_n\}$  runs through all subsets of  $\{1, \dots, n\}$  containing exactly  $m$  elements.

## Beginning of the proof of the lemma:

Let  $(U, u)$  be an oriented chart on  $M$ , diffeomorphic to  $Q_p^m$  with coordinates  $u^1 \geq 0, \dots, u^p \geq 0, u^{p+1}, \dots, u^m$ , such that  $\mu_0|_U = du^1 \wedge \dots \wedge du^m$ . The distribution  $\hat{G}|_U \in D'(U^n)$  has support contained in the full diagonal

$\Delta(U) = \{(x, \dots, x) \in U^n : x \in U\}$  and is of finite order  $k$  since  $M$  is compact. By Thm. 2.3.5 of Hörmander 1983, the corresponding multilinear form  $G$  can be written as

$$G(f_1, \dots, f_n) = \sum_{|\alpha_1| + \dots + |\alpha_{n-1}| \leq k} \langle A_{\alpha_1, \dots, \alpha_{n-1}}, \partial^{\alpha_1} f_1 \dots \partial^{\alpha_{n-1}} f_{n-1} \cdot f_n \rangle,$$

with multi-indices  $\alpha_j = (\alpha_{j,1}, \dots, \alpha_{j,m})$  and unique distributions  $A_{\alpha_1, \dots, \alpha_{n-1}} \in D'(U)$  of order  $k - |\alpha_1| - \dots - |\alpha_{n-1}|$ .

# End of the proof of the Main Theorem

Let  $\hat{G}$  be an invariant distribution in  $\mathcal{D}'(M^n)$  and let  $k < n/2$ . Let  $\{1, \dots, n\} = \{i_1, \dots, i_k\} \sqcup \{j_1, \dots, j_{n-k}\}$  be a partition into a disjoint union.

Without loss, let  $\{i_1, \dots, i_k\} = \{1, \dots, k\}$ . Let  $(x_1, \dots, x_n) \in M^n$  be such that no  $x_i$  for  $1 \leq i \leq k$  equals any of the  $x_j$  with  $k < j$ . Choose open neighborhoods  $U_{x_\ell}$  of  $x_\ell$  in  $M$  for all  $\ell$  such that each  $\overline{U_{x_i}}$  with  $i \leq k$  is disjoint from any  $\overline{U_{x_j}}$  with  $k < j$ . For smooth functions  $f_\ell$  with support in  $U_{x_\ell}$  for all  $\ell$ , we have that for  $i \leq k$  all functions  $f_i$  vanish on  $\bigcap_{j=1}^k (M \setminus U_{x_j})$ , thus

$\mathcal{L}_{X^{\text{diag}}}(\hat{G}(f_1, \dots, f_k))|(\bigcap_{j=1}^k (M \setminus U_{x_j}))^{n-k} = 0$  for all  $X \in \mathfrak{X}_{\text{diag}}(M, \partial M, \mu_0)$ .

For  $k < j$  we have  $\text{supp}(f_j) \subset U_{x_j} \subset \bigcap_{i=1}^k (M \setminus U_{x_i})$ . Consider  $f_1, \dots, f_k$  as fixed. Using induction on  $n$  and replacing  $M$  by the submanifold (non-compact!)  $\bigcap_{i=1}^k (M \setminus U_{x_i})$  we may assume that the main theorem is already true for

$$\check{G}_c(f_1, \dots, f_k) \Big| \left( \bigcap_{j=1}^k (M \setminus U_{x_j}) \right)^{n-k}$$

so that

$$\begin{aligned} \check{G}_c(f_1, \dots, f_k)(f_{k+1}, \dots, f_n) &= C_0(f_1, \dots, f_k) \int f_{k+1} \dots f_n \mu_0 \\ &+ \sum_{i=k+1}^n C_i(f_1, \dots, f_k) \int_M \alpha_i \cdot \int_M f_{k+1} \dots \widehat{f}_i \dots f_n \mu_0 \\ &+ \sum_{k < i < j}^n C_{ij}(f_1, \dots, f_k) \int_M f_i f_j \mu_0 \cdot \int_M f_{k+1} \dots \widehat{f}_i \dots \widehat{f}_j \dots f_n \mu \\ &+ \dots \\ &+ C_{12\dots n}(f_1, \dots, f_k) \int_M f_{k+1} \mu_0 \cdots \int_M f_n \mu. \end{aligned}$$

Now all the expressions  $C(f_1, \dots, f_k)$  are again invariant, and we can subject it also to the induction hypothesis. All the resulting multilinear operators are defined on the whole of  $M$ . If we subtract them from the original  $\hat{G}_C$ , the resulting distribution has support in the set of all  $(x_1, \dots, x_n) \in M^n$  such that  $x_{i_k} = x_{j_{\ell(k)}}$  for an injective mapping  $\ell : \{1, \dots, k\} \rightarrow \{1, \dots, n - k\}$ .

Finally we end up with a distribution with support on the full diagonal  $\{(x, \dots, x) : x \in M\} \subset M^n$  whose form is determined by the last lemma. □

Thank you for listening.